

ENUMERATION OF SPANNING SUBGRAPHS WITH DEGREE CONSTRAINTS

DAVID G. WAGNER

ABSTRACT. For a finite undirected multigraph $G = (V, E)$ and functions $f, g : V \rightarrow \mathbb{N}$, let $N_f^g(G, j)$ denote the number of (f, g) -factors of G with exactly j edges. The Heilmann-Lieb Theorem implies that $\sum_j N_0^1(G, j)t^j$ is a polynomial with only real (negative) zeros, and hence that the sequence $N_0^1(G, j)$ is strictly logarithmically concave. Separate generalizations of this theorem were obtained by Ruelle and by the author. We unify, simplify, and generalize these results by means of the Grace-Szegö-Walsh Coincidence Theorem.

1. INTRODUCTION.

By a *graph* $G = (V, E)$ we mean a finite undirected multigraph. We identify spanning subgraphs of G with subsets $H \subseteq E$ of edges, and let $\deg(H, v)$ denote the degree of $v \in V$ in the subgraph (V, H) . The *degree vector* of H is the function $\deg(H) : V \rightarrow \mathbb{N}$ given by $\deg(H)(v) := \deg(H, v)$ for all $v \in V$. Given functions $f, g : V \rightarrow \mathbb{N}$, an (f, g) -*factor* is a spanning subgraph H such that $f \leq \deg(H) \leq g$, in which the inequalities represent the coordinatewise partial order on $\mathbb{N}^V$. It is convenient to let natural numbers stand for constant functions on V , so that, for example, a $(0, 1)$ -factor of G is a matching in G . Let $N_f^g(G; j)$ denote the number of (f, g) -factors of G with exactly j edges. We are concerned here with obtaining inequalities among these numbers, and among weighted analogues of them.

This investigation is motivated by the following conjecture.

Conjecture 1. *For any graph $G = (V, E)$ and functions $f, g : V \rightarrow \mathbb{N}$, the sequence $\{N_f^g(G; j)\}$ is logarithmically concave: for all j ,*

$$N_f^g(G; j)^2 \geq N_f^g(G; j-1)N_f^g(G; j+1).$$

1991 *Mathematics Subject Classification.* 05A20; 05C30, 26C10, 30C15.

Key words and phrases. graph factor, Heilmann-Lieb theorem, Grace-Szegö-Walsh theorem, half-plane property, Hurwitz stability, logarithmic concavity.

Research supported by the Natural Sciences and Engineering Research Council of Canada under operating grant OGP0105392.

The evidence for this is admittedly meagre. The results reported here – and the amount of “slack” in their derivation – provide at least some heuristic support for the conjecture.

Conjecture 1 extrapolates from several results in the literature. The prototype is the famous (univariate version of a) theorem of Heilmann and Lieb [6] (see also Theorem 10.1 of [3]).

Theorem 2 (Heilmann–Lieb). *For any graph G , the polynomial $\sum_j N_0^1(G; j)t^j$ has only real (strictly negative) zeros.*

Newton’s Inequalities (Proposition 12 below) then imply that the sequence of coefficients is *strictly logarithmically concave*: if $N(j) > 0$ then $N(j)^2 > N(j-1)N(j+1)$.

More generally, we have the following (Theorem 3.3 of [14]).

Theorem 3. *For any graph $G = (V, E)$ and functions $f, g : V \rightarrow \mathbb{N}$ such that $f \leq g \leq f+1$, the polynomial $\sum_j N_f^g(G; j)t^j$ has only real (nonpositive) zeros.*

Ruelle [10] proves a result which relaxes the hypothesis $f \leq g \leq f+1$.

Theorem 4 (Ruelle). *For a graph G of maximum degree $\Delta \geq 2$, every zero of the polynomial $\sum_j N_0^2(G; j)t^j$ has real part less than or equal to $-2/\Delta(\Delta-1)^2$.*

A univariate polynomial for which all zeros have negative real part is said to be *Hurwitz stable*. This condition implies the inequalities $N(j)N(j+1) \geq N(j-1)N(j+2)$ among the coefficients (see Proposition 15 below.)

Another theorem of Ruelle [10] involves a weighted version of the numbers $N_0^2(G; j)$. Fix a sequence of nonnegative real numbers $\mathbf{u} := \{u_0, u_1, u_2, \dots\}$ (called *fugacities*), and for each $H \subseteq E$ let

$$\mathbf{u}_{\deg(H)} := \prod_{v \in V} u_{\deg(H, v)}.$$

For each natural number j let

$$N(G; \mathbf{u}, j) := \sum_{H \subseteq E: \#H=j} \mathbf{u}_{\deg(H)}.$$

For example, when $u_0 = u_1 = 1$ and $u_k = 0$ for $k \geq 2$ these are the numbers $N_0^1(G; j)$. Similarly, when $u_0 = u_1 = u_2 = 1$ and $u_k = 0$ for $k \geq 3$ these are the numbers $N_0^2(G; j)$. (More generally, if the functions f, g are not constant on V then the numbers $N_f^g(G; j)$ can be expressed in this way only if the fugacities are *anisotropic*: that is, they are allowed to vary from one vertex to another. Our main result is

anisotropic but for the purposes of this introduction the isotropic case above will suffice.)

Theorem 5 (Ruelle). *Let G be a graph with maximum degree $\Delta \geq 1$. Then for fugacities $\mathbf{u}$ satisfying $u_0 = u_2 = 1$, $u_1 \geq \sqrt{2 - 2/\Delta}$, and $u_k = 0$ for $k \geq 3$, the polynomial $\sum_j N(G; \mathbf{u}, j)t^j$ has only real (strictly negative) zeros.*

We prove the following result in Section 3. For a positive integer D and a sequence of fugacities $\mathbf{u}$, define the generating function

$$\Gamma(D, \mathbf{u}, y) := \sum_{k=0}^D \binom{D}{k} u_k y^k.$$

Proposition 6. *Let G be a graph with maximum degree Δ , and let $D \geq \Delta$ be a positive integer. Let $\mathbf{u}$ be a sequence of fugacities such that $\Gamma(D, \mathbf{u}, y)$ is a polynomial with only real nonpositive zeros. Then the polynomial $\sum_j N(G; \mathbf{u}, j)t^j$ has only real nonpositive zeros.*

In fact, Proposition 6 can be generalized in two directions – the fugacities can be replaced by an anisotropic set $\{\mathbf{u}^{(v)} : v \in V\}$ of fugacities, and the zeros of each $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ can be permitted to lie in a sector centered on the negative real axis (Corollary 19). The conclusion is then correspondingly weakened, but often allows the deduction of inequalities among the coefficients $\{N(G; \{\mathbf{u}^{(v)}\}, j)\}$. Proposition 6 itself implies both Theorems 2 and 5, as is easily verified. As we shall see, Corollary 19 also implies both Theorem 3 and a wide generalization of a slight weakening of Theorem 4. Therefore, Corollary 19 manages to unify all of the results presented in this introduction. Moreover, its proof is quite straightforward. Theorem 26 is an application of Corollary 19 which establishes a weak form of Conjecture 1.

Ruelle has a second paper [11] on this subject. His technique uses not only Grace’s theorem but also “Asano contraction” and some intricate geometry of polynomial zeros. He considers isotropic zero/one fugacities such that $u_k = 1$ if and only if $k \in S$, where S is among the following the sets: $\{0, 1\}$, $\{0, 1, 2\}$, $\{0, 2\}$, $\{0, 2, 4\}$, $\{k \text{ even}\}$, $\{k < \Delta\}$, $\{k \geq 1\}$. In these cases Ruelle produces more precise information about the location of zeros of $\sum_j N(G; \mathbf{u}, j)t^j$ than can be obtained by our method. (On the other hand, our method is very easily applied to a wide class of vertex degree restrictions.) Moreover, Ruelle considers factors of directed graphs in which the indegrees and outdegrees are subject to separate restrictions. We indicate briefly at the end how our method can also be extended to the case of directed graphs. Perhaps

some elaboration of Ruelle's ideas and our own could lead to further progress towards Conjecture 1.

I thank Alan Sokal for stimulating my interest in the techniques of this paper, and in particular for showing me the usefulness of the Grace–Szegő–Walsh Coincidence Theorem.

2. PRELIMINARIES.

As indicated in the introduction, we proceed by locating the zeros of polynomials within certain prescribed sectors and circles. In this section we first collect the necessary tools, and then explain the implications for coefficient inequalities.

Let $F(\mathbf{z})$ be a polynomial in complex variables $\mathbf{z} := \{z_v : v \in V\}$. For an open subset $\mathcal{A} \subset \mathbb{C}$, we say that F is $\mathcal{A}$ -nonvanishing if either $F \equiv 0$, or $z_v \in \mathcal{A}$ for all $v \in V$ implies that $F(\mathbf{z}) \neq 0$. If $F \not\equiv 0$ then we say that F is *strictly* $\mathcal{A}$ -nonvanishing.

The following lemma is obvious, and its proof is omitted.

Lemma 7. *Let $F(\mathbf{z})$ be $\mathcal{A}$ -nonvanishing, and let $S \subseteq V$. Let $\tilde{z}_v = z_v$ if $v \notin S$, and let $\tilde{z}_v = z$ if $v \in S$. Then $F(\tilde{\mathbf{z}})$ is $\mathcal{A}$ -nonvanishing.*

Lemma 8. *Let $F_\rho(\mathbf{z})$ be a family of strictly $\mathcal{A}$ -nonvanishing polynomials indexed by positive real numbers $\rho \in (0, \varepsilon)$. Assume that the limit $F(\mathbf{z}) := \lim_{\rho \rightarrow 0} F_\rho(\mathbf{z})$ exists. Then F is $\mathcal{A}$ -nonvanishing.*

Proof. Each F_ρ is analytic and nonvanishing on the subset $\mathcal{A}^V$ of $\mathbb{C}^V$. Since these functions are polynomials, the convergence to F is uniform on compact subsets of $\mathbb{C}^V$. By Hurwitz's Theorem, either F is identically zero or F is nonvanishing on $\mathcal{A}^V$ as well. $\square$

Lemma 9. *Let $F(\mathbf{z})$ be $\mathcal{A}$ -nonvanishing, and let $w \in V$. If z_w is fixed at a complex value ξ_0 in the closure of $\mathcal{A}$, then the resulting polynomial in the variables $\{z_v : v \in V \setminus \{w\}\}$ is $\mathcal{A}$ -nonvanishing.*

Proof. Let $\xi : (0, \varepsilon) \rightarrow \mathcal{A}$ be a continuous function such that $\lim_{\rho \rightarrow 0} \xi(\rho) = \xi_0$. For any positive value of ρ , the specialization $z_w = \xi(\rho)$ results in a polynomial F_ρ that is $\mathcal{A}$ -nonvanishing in the variables $\{z_v : v \in V \setminus \{w\}\}$. The result follows from Lemma 8. $\square$

Of particular importance here are the following subsets of $\mathbb{C}$:

(i) For $0 < \theta \leq \pi$, the open sector

$$\mathcal{S}[\theta] := \{z \in \mathbb{C} : z \neq 0 \text{ and } |\arg(z)| < \theta\}.$$

(For $z \neq 0$ we use the value of the argument in the range $-\pi < \arg(z) \leq \pi$.) For the open right half-plane $\mathcal{H} := \mathcal{S}[\pi/2]$ an $\mathcal{H}$ -vanishing polynomial is also said to be *Hurwitz quasi-stable* or have the *half-plane*

property. Notice that if F is $\mathcal{S}[\pi]$ -nonvanishing and $F(\mathbf{z}) = 0$ then at least one of the complex numbers z_v must be a nonpositive real number. In particular, a univariate polynomial is $\mathcal{S}[\pi]$ -nonvanishing if and only if it has only real nonpositive zeros.

(ii) The open unit disc $\mathcal{D} := \{z \in \mathbb{C} : |z| < 1\}$.

(iii) The open exterior of the unit disc $\mathcal{E} := \{z \in \mathbb{C} : |z| > 1\}$.

For a positive real number $\kappa > 0$, we let $\kappa\mathcal{D} := \{z \in \mathbb{C} : |z| < \kappa\}$ and $\kappa\mathcal{E} := \{z \in \mathbb{C} : |z| > \kappa\}$.

The sets $\mathcal{H}$, $\kappa\mathcal{D}$, and $\kappa\mathcal{E}$ are examples of *circular regions*, in that they are bounded by either circles or straight lines. For such regions, the Grace–Szegő–Walsh Coincidence Theorem can be extremely useful. For a polynomial $F(\mathbf{z})$, a vertex $w \in V$, and an integer D greater than or equal to the maximum degree to which z_w occurs in F , the D -th *polarization of z_w in F* is the polynomial $P_w^D F(\mathbf{z})$ defined as follows. Introduce new variables $\{z_{w1}, \dots, z_{wD}\}$ and let $e_k(z_{w1}, \dots, z_{wD})$ denote the k -th elementary symmetric function of $\{z_{w1}, \dots, z_{wD}\}$. Then $P_w^D F(\mathbf{z})$ is obtained from $F(\mathbf{z})$ by applying the linear transformation defined by $z_w^k \mapsto \binom{D}{k}^{-1} e_k(z_{w1}, \dots, z_{wD})$ and linear extension.

Proposition 10 (Grace–Szegő–Walsh). *Let $\mathcal{A}$ be a circular region, let $F(\mathbf{z})$ be $\mathcal{A}$ -nonvanishing, let $w \in V$, and let D be an integer greater than or equal to the maximum degree $d(w)$ to which z_w occurs in $F(\mathbf{z})$. If either $D = d(w)$ or $\mathcal{A}$ is convex then $P_w^D F(\mathbf{z})$ is also $\mathcal{A}$ -nonvanishing.*

Theorem 15.4 of Marden [9] provides a proof for the case $D = d(w)$. The theorem also holds for $D > d(w)$ with the additional hypothesis that $\mathcal{A}$ is convex, as explained in Theorem 2.12 of [3].

Proposition 11 (Takagi, Weisner). *Let $p(y) = \sum_{k=0}^n a_k y^k$ and $q(y) = \sum_{k=0}^n b_k y^k$ be real polynomials of degree at most n . Assume that $p(y)$ is $\mathcal{S}[\pi]$ -nonvanishing, and that $q(y)$ is $\mathcal{S}[\theta]$ -nonvanishing for some $\theta > \pi/2$. Then each of the following polynomials is $\mathcal{S}[\theta]$ -nonvanishing.*

- (a) $\sum_{k=0}^n a_k b_k y^k$;
- (b) $\sum_{k=0}^n k! a_k b_k y^k$;
- (c) $\sum_{k=0}^n k!(n-k)! a_k b_k y^k$.

For a proof, see Takagi [12], Weisner [15], or the exercises in Section 16 of Marden [9]. Related results can be found in [4, 13].

Location of zeros of generating functions implies combinatorial inequalities *via* the following results.

Proposition 12 (Newton's Inequalities). *Let $F(t) = \sum_{j=0}^d N(j)t^j$ be a univariate polynomial with real coefficients and only real zeros. Then*

$$\frac{N(j)^2}{\binom{d}{j}^2} \geq \frac{N(j-1)}{\binom{d}{j-1}} \cdot \frac{N(j+1)}{\binom{d}{j+1}}$$

for all $1 \leq j \leq d-1$. In particular, $N(j)^2 > N(j-1)N(j+1)$ whenever $N(j) > 0$.

See inequality (51) of Hardy–Littlewood–Pólya [5] for a proof.

Proposition 13 (Aissen–Schoenberg–Whitney). *Let $F(t) = \sum_{j=0}^d N(j)t^j$ be a univariate polynomial with nonnegative coefficients. Then $F(t)$ has only real nonpositive zeros if and only if every finite minor of the Toeplitz matrix*

$$\begin{bmatrix} N(0) & N(1) & N(2) & N(3) & \cdots \\ 0 & N(0) & N(1) & N(2) & \cdots \\ 0 & 0 & N(0) & N(1) & \cdots \\ 0 & 0 & 0 & N(0) & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}$$

is nonnegative. In particular, in this case then

$$N(j)^2 \geq N(j-1)N(j+1)$$

for all $1 \leq j \leq d-1$.

See [1] or Chapter 8 of Karlin [7] for a proof.

Proposition 14 (Folklore). *Let $F(t) = \sum_{j=0}^d N(j)t^j$ be a univariate polynomial with real coefficients. If F is $\mathcal{S}[2\pi/3]$ -nonvanishing then $N(j)^2 \geq N(j-1)N(j+1)$ for all $1 \leq j \leq d-1$.*

Proof. Factor $F(t)$ over the reals, and proceed by induction on the degree of $F(t)$, using the fact that if $p(t)$ and $q(t)$ are polynomials with logarithmically concave sequences of coefficients, then $p(t)q(t)$ also has a logarithmically concave sequence of coefficients. $\square$

Proposition 15 (Hurwitz). *Let $F(t) = \sum_{j=0}^d N(j)t^j$ be a univariate polynomial with nonnegative coefficients. Then $F(t)$ is $\mathcal{H}$ -nonvanishing*

if and only if every minor of the Hurwitz matrix

$$\begin{bmatrix} N(1) & N(3) & N(5) & \cdots & & 0 \\ N(0) & N(2) & N(4) & \cdots & & 0 \\ 0 & N(1) & N(3) & \cdots & & 0 \\ 0 & N(0) & N(2) & \cdots & & 0 \\ \vdots & & & \ddots & & \vdots \\ 0 & & \cdots & N(d-3) & N(d-1) & 0 \\ 0 & 0 & \cdots & & N(d-2) & N(d) \end{bmatrix}$$

is nonnegative. In particular, in this case then

$$N(j)N(j+1) \geq N(j-1)N(j+2)$$

for all $1 \leq j \leq d-2$, and

$$N(j)^2 \geq N(j-2)N(j+2)$$

for all $2 \leq j \leq d-2$.

See Asner [2] or Kemperman [8] for a proof.

3. RESULTS.

For a graph $G = (V, E)$, let $\boldsymbol{\lambda} := \{\lambda_e : e \in E\}$ be positive real constants indexed by the edges of G , and let $\mathbf{z} := \{z_v : v \in V\}$ be complex variables indexed by the vertices of G . We indicate that the ends of $e \in E$ are the vertices v and w by writing $vew \in E$ (of course, $v = w$ is possible). Notice that

$$F(G; \boldsymbol{\lambda}, \mathbf{z}) := \prod_{vew \in E} (1 + \lambda_e z_v z_w) = \sum_{H \subseteq E} \boldsymbol{\lambda}_H \mathbf{z}^{\deg(H)}$$

is a weighted multivariate generating function for all spanning subgraphs of G , in which $\boldsymbol{\lambda}_H := \prod_{e \in H} \lambda_e$.

Proposition 16. *Let $G = (V, E)$ be a graph, and fix positive constants $\boldsymbol{\lambda} = \{\lambda_e : e \in E\}$ such that $\lambda_{\min} \leq \lambda_e \leq \lambda_{\max}$ for all $e \in E$. Then*

- (a) $F(G; \boldsymbol{\lambda}, \mathbf{z})$ is $\mathcal{H}$ -nonvanishing.
- (b) $F(G; \boldsymbol{\lambda}, \mathbf{z})$ is $\lambda_{\max}^{-1/2} \mathcal{D}$ -nonvanishing.
- (c) $F(G; \boldsymbol{\lambda}, \mathbf{z})$ is $\lambda_{\min}^{-1/2} \mathcal{E}$ -nonvanishing.

Proof. In each case, each factor $1 + \lambda_e z_v z_w$ in the product $F(G; \boldsymbol{\lambda}, \mathbf{z})$ is seen to be nonvanishing in the appropriate region, from which the result follows. $\square$

The main theorem of this paper is as follows. Let $G = (V, E)$ be a graph, and for each vertex $v \in V$ fix a sequence of nonnegative fugacities $\mathbf{u}^{(v)} := \{u_0^{(v)}, u_1^{(v)}, u_2^{(v)}, \dots\}$. For $H \subseteq E$, let

$$\mathbf{u}_{\deg(H)} := \prod_{v \in V} u_{\deg(H, v)}^{(v)}.$$

Theorem 17. *Let $G = (V, E)$ be a graph, and let $D : V \rightarrow \mathbb{N}$ be such that $\deg(G) \leq D$. For each $v \in V$, fix a sequence $\mathbf{u}^{(v)}$ of nonnegative fugacities with generating function $\Gamma_v(y) := \Gamma(D(v), \mathbf{u}^{(v)}, y)$. Consider the polynomial $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z}) := \sum_{H \subseteq E} \mathbf{u}_{\deg(H)} \mathbf{z}^{\deg(H)}$.*

- (a) *Fix $0 \leq \alpha < \pi/2$. If $\Gamma_v(y)$ is $\mathcal{S}[\pi - \alpha]$ -nonvanishing for all $v \in V$ then $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z})$ is $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing.*
- (b) *Fix $\kappa > 0$. If $\Gamma_v(y)$ is $\kappa\mathcal{D}$ -nonvanishing for all $v \in V$ then $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z})$ is $\kappa\mathcal{D}$ -nonvanishing.*
- (c) *Fix $\kappa > 0$. If $D = \deg(G)$, $u_{D(v)}^{(v)} \neq 0$, and $\Gamma_v(y)$ is $\kappa\mathcal{E}$ -nonvanishing for all $v \in V$ then $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z})$ is $\kappa\mathcal{E}$ -nonvanishing.*

Proof. We begin with part (a). First, we prove the special case in which each $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ is $\mathcal{S}[\pi - \alpha]$ -nonvanishing and also has a nonzero constant term. Afterward, the general case will be obtained by a limiting argument. Thus, in this special case we have, for each $v \in V$,

$$\Gamma_v(y) = \sum_{k=0}^{D(v)} \binom{D(v)}{k} u_k^{(v)} y^k = u_0^{(v)} \prod_{i=1}^{D(v)} (1 + \xi_{vi} y),$$

for some complex numbers $\{\xi_{vi}\}$ such that either $\xi_{vi} = 0$ or $|\arg(\xi_{vi})| \leq \alpha$ for all $v \in V$ and $1 \leq i \leq D(v)$.

Let $F(G; \mathbf{z})$ be the polynomial of Proposition 16 in which all $\lambda_e = 1$. This polynomial is $\mathcal{H}$ -nonvanishing. For each $v \in V$, perform the $D(v)$ -fold polarization of the variable z_v in $F(G; \mathbf{z})$. Let us denote the resulting polynomial by $P^D F(G; \{z_{vi}\})$. Since $D(v) \geq \deg(G, v)$ for each $v \in V$, repeated application of Proposition 10 implies that $P^D F(G; \{z_{vi}\})$ is $\mathcal{H}$ -nonvanishing. Next, replace the variable z_{vi} by $\xi_{vi} z_{vi}$ for all $v \in V$ and $1 \leq i \leq D(v)$. Since either $\xi_{vi} = 0$ or $|\arg(\xi_{vi})| \leq \alpha$ for all v and i , and since $P^D F(G; \{z_{vi}\})$ is $\mathcal{H}$ -nonvanishing, it follows that the resulting polynomial $P^D F(G; \{\xi_{vi} z_{vi}\})$ is $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing. Finally, partially diagonalize the variables by making the substitutions $z_{vi} \mapsto z_v$ for all $v \in V$ and $1 \leq i \leq D(v)$. Repeated application of Lemma 7 shows that the resulting polynomial $P^D F(G; \{\xi_{vi} z_v\})$ is still $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing. Consider the effect

of these operations on z_v^k in $F(G; \mathbf{z})$:

$$z_v^k \mapsto \binom{D(v)}{k}^{-1} e_k(\xi_{v1}z_v, \dots, \xi_{vD(v)}z_v) = u_k^{(v)} z_v^k,$$

since $e_k(\xi_{v1}, \dots, \xi_{vD(v)}) = \binom{D(v)}{k} u_k^{(v)}$. Therefore,

$$P^D F(G; \{\xi_{vi}z_v\}) = \sum_{H \subseteq E} \mathbf{u}_{\deg(H)} \mathbf{z}^{\deg(H)}$$

is $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing, completing the proof in this case.

We indicate how the general case of part (a) is derived from the special case above by relaxing the hypothesis at one vertex $w \in V$. Iteration of this argument for each vertex then yields part (a) as stated. Accordingly, assume that

$$\Gamma(D(w), \mathbf{u}^{(w)}, y) = u_m^{(w)} y^m \prod_{i=1}^{D(w)-m} (1 + \xi_{wi}y)$$

for some complex constants $\{\xi_{wi}\}$ with either $\xi_{wi} = 0$ or $|\arg(\xi_{wi})| \leq \alpha$. Fix a positive real number $\rho > 0$, and replace the fugacities $\mathbf{u}^{(w)}$ by those $\tilde{\mathbf{u}}^{(w)}$ with generating function

$$\tilde{\Gamma}_w(y) = (\rho + y)^m \prod_{i=1}^{D(w)-m} (1 + \xi_{wi}y).$$

One easily verifies that as $\rho \rightarrow 0$,

$$\tilde{u}_i^{(w)} \rightarrow \begin{cases} 0 & \text{if } 0 \leq i < m, \\ u_i^{(w)} & \text{if } m \leq i \leq D(w). \end{cases}$$

The special case above shows that for any positive value of $\rho > 0$, $\sum_{H \subseteq E} \tilde{\mathbf{u}}_{\deg(H)} \mathbf{z}^{\deg(H)}$ is $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing. By Lemma 8, it follows that $\sum_{H \subseteq E} \mathbf{u}_{\deg(H)} \mathbf{z}^{\deg(H)}$ is also $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing, as required.

For part (b), the hypothesis implies that every zero of every $\Gamma_v(y)$ has modulus at least κ . In particular, for each $v \in V$ we have $u_0^{(v)} > 0$ and

$$\Gamma_v(y) = u_0^{(v)} \prod_{i=1}^{D(v)} (1 + \xi_{vi}y)$$

for some complex numbers $\{\xi_{vi}\}$ such that $|\xi_{vi}| \leq 1/\kappa$ for all $1 \leq i \leq D(v)$.

Propositions 16 and 10 imply that $P^D F(G; \{z_{vi}\})$ is $\mathcal{D}$ -nonvanishing. Replacing each z_{vi} by $\xi_{vi}z_{vi}$, the resulting polynomial $P^D F(G; \{\xi_{vi}z_{vi}\})$ is $\kappa\mathcal{D}$ -nonvanishing. Upon making the substitutions $z_{vi} \mapsto z_v$ for all

$v \in V$ and $1 \leq i \leq D(v)$, Lemma 7 implies that $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z})$ is $\kappa\mathcal{D}$ -nonvanishing.

For part (c) the hypothesis implies that every zero of every $\Gamma_v(y)$ has modulus at most κ , and that $u_{D(v)}^{(v)} > 0$. For each $v \in V$ let $m(v)$ be the multiplicity of 0 as a root of $\Gamma_v(y)$; we may write

$$\Gamma_v(y) = \lim_{\rho \rightarrow 0} \rho^{m(v)} u_m^{(v)} (1 + y/\rho)^{m(v)} \prod_{i=1}^{D(v)-m(v)} (1 + \xi_{vi}y)$$

for some complex numbers $\{\xi_{vi}\}$ such that $|\xi_{vi}| \geq 1/\kappa$ for all $1 \leq i \leq D(v) - m(v)$. Let $\xi_{vi} = 1/\rho$ for $D(v) - m(v) + 1 \leq i \leq D(v)$.

Propositions 16 and 10 imply that $P^D F(G; \{z_{vi}\})$ is $\mathcal{E}$ -nonvanishing. For any positive $\rho \leq \kappa$, upon replacing each z_{vi} by $\xi_{vi}z_{vi}$, the resulting polynomial $P^D F(G; \{\xi_{vi}z_{vi}\})$ is $\kappa\mathcal{E}$ -nonvanishing. The substitutions $z_{vi} \mapsto z_v$ for all $v \in V$ and $1 \leq i \leq D(v)$, and Lemma 7 imply that $P^D F(G; \{\xi_{vi}z_v\})$ is $\kappa\mathcal{D}$ -nonvanishing. With $M := \sum_{v \in V} m(v)$, the limit of $\rho^M P^D F(G; \{\xi_{vi}z_v\})$ as $\rho \rightarrow 0$ is $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z})$. Lemma 8 then shows that $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z})$ is $\kappa\mathcal{E}$ -nonvanishing, as desired. $\square$

We can specialize Theorem 17 immediately to obtain a multivariate generalization of Theorem 3.

Corollary 18. *For any graph $G = (V, E)$ and functions $f, g : V \rightarrow \mathbb{N}$ such that $f \leq g \leq f + 1$, the polynomial*

$$\sum_{H \subseteq E: f \leq \deg(H) \leq g} \mathbf{z}^{\deg(H)}$$

is $\mathcal{H}$ -nonvanishing.

Proof. Apply Theorem 17(a) by taking $D = \deg(G)$, $\alpha = 0$, and

$$\Gamma(D(v), \mathbf{u}^{(v)}, y) = \sum_{k=f(v)}^{g(v)} \binom{D(v)}{k} y^k$$

for each $v \in V$. $\square$

The case $f \equiv 0$ and $g \equiv 1$ of Corollary 18 is the multivariate version of the Heilmann–Lieb theorem [6].

With the notation of Theorem 17, for $j \in \mathbb{N}$ let

$$N(G; \{\mathbf{u}^{(v)}\}, j) := \sum_{H \subseteq E: \#H=j} \mathbf{u}_{\deg(H)}.$$

Corollary 19. *Let $G = (V, E)$ be a graph, and let $D : V \rightarrow \mathbb{N}$ be such that $\deg(G) \leq D$. Fix $0 \leq \alpha < \pi/2$. For each $v \in V$, let $\mathbf{u}^{(v)}$ be a*

sequence of nonnegative fugacities such that $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ is $\mathcal{S}[\pi - \alpha]$ -nonvanishing. Then the univariate polynomial $\sum_j N(G; \{\mathbf{u}^{(v)}\}, j)t^j$ is $\mathcal{S}[\pi - 2\alpha]$ -nonvanishing.

Proof. By Theorem 17, the polynomial $\sum_{H \subseteq E} \mathbf{u}_{\deg(H)} \mathbf{z}^{\deg(H)}$ is $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing. Fully diagonalize this, by making the substitutions $z_v \mapsto z$ for all $v \in V$. By Lemma 7, the resulting polynomial $\sum_j N(G; \{\mathbf{u}^{(v)}\}, j)z^{2j}$ is also $\mathcal{S}[\pi/2 - \alpha]$ -nonvanishing. Since every complex value t with $|\arg(t)| < \pi - 2\alpha$ has a square root z with $|\arg(z)| < \pi/2 - \alpha$, it follows that the polynomial $\sum_j N(G; \{\mathbf{u}^{(v)}\}, j)t^j$ is $\mathcal{S}[\pi - 2\alpha]$ -nonvanishing. $\square$

Corollary 20. *Let $G = (V, E)$ be a graph, let $D : V \rightarrow \mathbb{N}$ be such that $\deg(G) \leq D$, and to every $v \in V$ assign nonnegative fugacities $\mathbf{u}^{(v)}$ such that*

$$Q(\mathbf{u}^{(v)}, y) := \sum_{k=0}^{D(v)} u_k^{(v)} \frac{y^k}{k!}$$

is $\mathcal{S}[\pi - \alpha]$ -nonvanishing, with $0 \leq \alpha < \pi$. Then the polynomial $\sum_j N(G; \{\mathbf{u}^{(v)}\}, j)t^j$ is $(\pi - 2\alpha)$ -nonvanishing.

Proof. For each $v \in V$, Proposition 11(b) with $p(y) = (1 + y)^{D(v)}$ and $q(y) = Q(\mathbf{u}^{(v)}, y)$ shows that the generating function $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ is $\mathcal{S}[\pi - \alpha]$ -nonvanishing. The result follows immediately from Corollary 19. $\square$

Theorem 3.2 of [14] is the case $\alpha = 0$ of Corollary 20. This in turn implies Theorems 2 and 3 above.

Corollary 19 has the following simple consequence. As grist for the mill we need to locate the zeros of some quadratic and cubic polynomials within certain sectors. The proof of Lemma 21 is a routine calculation, which is omitted.

Lemma 21. *Let $1 \leq k \leq D - 1$ be integers, $\beta \geq 0$, let*

$$\Gamma(y) = \binom{D}{k-1} y^{k-1} + \beta \binom{D}{k} y^k + \binom{D}{k+1} y^{k+1},$$

and let $R = k(D - k)/(k + 1)(D - k + 1)$.

- (a) *If $\beta \geq \sqrt{2R}$ then $\Gamma(y)$ is strictly $\mathcal{S}[3\pi/4]$ -nonvanishing.*
- (b) *If $\beta \geq \sqrt{3R}$ then $\Gamma(y)$ is strictly $\mathcal{S}[5\pi/6]$ -nonvanishing.*
- (c) *If $\beta \geq 2\sqrt{R}$ then $\Gamma(y)$ is strictly $\mathcal{S}[\pi]$ -nonvanishing.*

Lemma 22. *Let $1 \leq k \leq D - 2$ be integers, let $\mu \geq 0$, and let*

$$\Gamma(y) = \binom{D}{k-1} y^{k-1} + \mu \binom{D}{k} y^k + \mu \binom{D}{k+1} y^{k+1} + \binom{D}{k+2} y^{k+2}.$$

- (a) If $\mu \geq 1 + \sqrt{2}$ then $\Gamma(y)$ is strictly $\mathcal{S}[3\pi/4]$ -nonvanishing.
- (b) If $\mu \geq 1 + \sqrt{3}$ then $\Gamma(y)$ is strictly $\mathcal{S}[5\pi/6]$ -nonvanishing.
- (c) If $\mu \geq 3$ then $\Gamma(y)$ is strictly $\mathcal{S}[\pi]$ -nonvanishing.

Proof. Let $p(y) = (1+y)^D$ and $q(y) = y^{k-1}(1+\mu y+\mu y^2+y^3)$. Hypothesis (a) implies that $q(y)$ is $\mathcal{S}[3\pi/4]$ -nonvanishing, hypothesis (b) implies that $q(y)$ is $\mathcal{S}[5\pi/6]$ -nonvanishing, and hypothesis (c) implies that $q(y)$ is $\mathcal{S}[\pi]$ -nonvanishing. The conclusion now follows from Proposition 11(a). $\square$

Lemma 22 ignores some “finite D ” effects that can be significant, especially for small D . The general case is quite complicated, but Lemma 23 is indicative of the possible improvement. Again, the elementary calculation is omitted.

Lemma 23. *Let $p \geq 1$ be an integer, $D = 2p + 1$, $\mu \geq 0$, and let*

$$\Gamma(y) = \binom{2p+1}{p-1} y^{p-1} + \mu \binom{2p+1}{p} y^p + \mu \binom{2p+1}{p+1} y^{p+1} + \binom{2p+1}{p+2} y^{p+2}.$$

- (a) If $\mu \geq (1 + \sqrt{2})p/(p+2)$ then $\Gamma(y)$ is strictly $\mathcal{S}[3\pi/4]$ -nonvanishing.
- (b) If $\mu \geq (1 + \sqrt{3})p/(p+2)$ then $\Gamma(y)$ is strictly $\mathcal{S}[5\pi/6]$ -nonvanishing.
- (c) If $\mu \geq 3p/(p+2)$ then $\Gamma(y)$ is strictly $\mathcal{S}[\pi]$ -nonvanishing.

Proposition 24. *Let $G = (V, E)$ be a graph, let $D : V \rightarrow \mathbb{N}$ be such that $\deg(G) \leq D$, and assign nonnegative fugacities $\mathbf{u}^{(v)}$ to each $v \in V$ so that each $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ has $r(v) \leq 4$ nonzero terms, of consecutive degrees. In each case below, assume that the generating functions for $v \in V$ with $r(v) \geq 3$ satisfy the given hypotheses.*

- (a) *Lemmas 21, 22, and 23, part (a): then $F(G; \{\mathbf{u}^{(v)}\}, t)$ is $\mathcal{H}$ -nonvanishing. In this case the inequalities of Proposition 15 hold for $\{N(G; \{\mathbf{u}^{(v)}\}, j)\}$.*
- (b) *Lemmas 21, 22, and 23, part (b): then $F(G; \{\mathbf{u}^{(v)}\}, t)$ is $\mathcal{S}[2\pi/3]$ -nonvanishing. In this case the inequalities of Proposition 14 hold for $\{N(G; \{\mathbf{u}^{(v)}\}, j)\}$.*
- (c) *Lemmas 21, 22, and 23, part (c): then $F(G; \{\mathbf{u}^{(v)}\}, t)$ has only real nonpositive zeros. In this case the inequalities of Propositions 12 and 13 hold for $\{N(G; \{\mathbf{u}^{(v)}\}, j)\}$.*

Proof. In part (a) every $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ is $\mathcal{S}[3\pi/4]$ -nonvanishing. In part (b) every $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ is $\mathcal{S}[5\pi/6]$ -nonvanishing. In part (c) every $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ is $\mathcal{S}[\pi]$ -nonvanishing. The result follows immediately from Corollary 19. $\square$

Of most interest combinatorially is the case in which all fugacities are either zero or one. All we obtain in this direction is the following rather limited result.

Proposition 25. *Let $G = (V, E)$ be a graph, and let $f, g : V \rightarrow \mathbb{N}$ be functions such that $f \leq g \leq f + 2$ and $g \leq \deg(G)$.*

- (a) *Then the polynomial $\sum_j N_f^g(G; j)t^j$ is $\mathcal{S}[\pi/3]$ -nonvanishing.*
- (b) *Assume furthermore that for every $v \in V$, either $g(v) \leq f(v) + 1$, or $f(v) = 0$, or $g(v) = \deg(G, v)$, or $\deg(G, v) \leq 5$. Then the polynomial $\sum_j N_f^g(G; j)t^j$ is $\mathcal{H}$ -nonvanishing. Thus the inequalities of Proposition 15 hold for $\{N_f^g(G; j)\}$.*

Proof. We apply Corollary 19 with $D = \deg(G)$ and fugacities given by

$$u_i^{(v)} := \begin{cases} 1 & \text{if } f(v) \leq i \leq g(v), \\ 0 & \text{otherwise,} \end{cases}$$

for all $v \in V$ and $i \in \mathbb{N}$. In this case, if $g(v) \leq f(v) + 1$ then $\Gamma_v(y) = \Gamma(D(v), \mathbf{u}^{(v)}, y)$ is $\mathcal{S}[\pi]$ -nonvanishing. If $f(v) + 1 = k = g(v) - 1$ then $1 \leq k \leq D(v) - 1$ and $\Gamma_v(y)$ has the form in Lemma 21 with $\beta = 1$. A short calculation shows that $\Gamma_v(y)$ is $\mathcal{S}[2\pi/3]$ -nonvanishing. Corollary 19 thus implies part (a). For part (b) one checks that $\Gamma_v(y)$ is $\mathcal{S}[3\pi/4]$ -nonvanishing if and only if either $k = 1$ or $k = D(v) - 1$ or $D(v) \leq 5$, and thus Corollary 19 implies part (b). $\square$

Our last application of Corollary 19 is a set of fugacities $\{\mathbf{u}^{(v)}\}$ weighting the (f, g) -factors of a graph that is sufficient to imply logarithmic concavity of the resulting numbers. This can be regarded as a (very) weak form of Conjecture 1.

Theorem 26. *Let $G = (V, E)$ be a graph, $D = \deg(G)$, and let $f, g : V \rightarrow \mathbb{N}$ with $f \leq g \leq D$. For each $v \in V$ let $g(v) - f(v) = 2a_v + b_v$ with $0 \leq b_v \leq 1$, and define*

$$q_v(y) = y^{f(v)}(1 + y)^{b_v}(1 + \sqrt{3}y + y^2)^{a_v}.$$

Assign nonnegative fugacities $\mathbf{u}^{(v)}$ to $v \in V$ so that $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ is the result of applying Proposition 11(a) to $(1 + y)^{D(v)}$ and $q_v(y)$. Then the polynomial $F(G; \{\mathbf{u}^{(v)}\}, t)$ is $\mathcal{S}[2\pi/3]$ -nonvanishing. Therefore, the coefficients $\{N(G; \{\mathbf{u}^{(v)}\}, j)\}$ are logarithmically concave.

Proof. By construction, each $q_v(y)$ is $\mathcal{S}[5\pi/6]$ -nonvanishing. By Proposition 11(a), the same is true for each $\Gamma(D(v), \mathbf{u}^{(v)}, y)$. The result follows immediately from Corollary 19 and Proposition 14. $\square$

Replacing the quadratic $1 + \sqrt{3}y + y^2$ in Theorem 26 by $1 + \sqrt{2}y + y^2$ gives a sufficient condition for $F(G; \{\mathbf{u}^{(v)}\}, t)$ to be $\mathcal{H}$ -nonvanishing. Using $1 + 2y + y^2$ instead gives a sufficient condition for $F(G; \{\mathbf{u}^{(v)}\}, t)$ to be $\mathcal{S}[\pi]$ -nonvanishing.

Parts (b) and (c) of Theorem 17 have the following consequence. For a graph $G = (V, E)$ and functions $f, g : V \rightarrow \mathbb{N}$, let

$$\binom{g}{f} := \prod_{v \in V} \binom{g(v)}{f(v)}.$$

Theorem 27. *Let $G = (V, E)$ be a graph.*

(a) *Let $D = \deg(G)$ and assign nonnegative fugacities $\mathbf{u}^{(v)}$ to each $v \in V$ so that $\Gamma(D(v), \mathbf{u}^{(v)}, y)$ has degree $D(v)$ and only zeros of unit modulus. Then every zero of $F(G; \{\mathbf{u}^{(v)}\}, t)$ has unit modulus.*

(b) *In particular, every zero of the polynomial*

$$\sum_{H \subseteq E} \binom{\deg(G)}{\deg(H)}^{-1} t^{\#H}$$

has unit modulus.

Proof. For part (a), the hypotheses of Theorem 17(b,c) are satisfied with $\kappa = 1$, so that $F(G; \{\mathbf{u}^{(v)}\}, \mathbf{z})$ is both $\mathcal{D}$ -nonvanishing and $\mathcal{E}$ -nonvanishing. After diagonalizing all the variables $z_v \mapsto t^{1/2}$, Lemma 7 yields the result.

Part (b) is the special case of part (a) in which the fugacities are $u_i^{(v)} = \binom{D(v)}{i}^{-1}$ for all $v \in V$ and $0 \leq i \leq D(v)$, so that the generating functions are $\Gamma(D(v), \mathbf{u}^{(v)}, y) = 1 + y + y^2 + \cdots + y^{D(v)} = (1 - y^{1+D(v)})/(1 - y)$. $\square$

4. DIRECTED GRAPHS.

We can rework the machinery of the previous section for directed graphs, as follows. Let $G = (V, E)$ be a directed graph. For a spanning directed subgraph $H \subseteq E$ of G , let $\text{outdeg}(H)$ and $\text{indeg}(H)$ denote the vectors of outdegrees and of indegrees in H , respectively. We use the notation $vw \in E$ to denote that e is an edge of G directed out of v and into w . Associate two sets of complex variables $\mathbf{z}' := \{z'_v : v \in V\}$ and $\mathbf{z}'' := \{z''_v : v \in V\}$ with the vertices of G . Fix positive real weights $\boldsymbol{\lambda} := \{\lambda_e : e \in E\}$ for the edges of G . The polynomial

$$F(G; \boldsymbol{\lambda}, \mathbf{z}', \mathbf{z}'') := \prod_{vw \in E} (1 + \lambda_e z'_v z''_w) = \sum_{H \subseteq E} \boldsymbol{\lambda}_H(\mathbf{z}')^{\text{outdeg}(H)} (\mathbf{z}'')^{\text{indeg}(H)}$$

is a weighted generating function for all spanning directed subgraphs of G . As in Proposition 16 this polynomial is $\mathcal{H}$ -nonvanishing, and if $\boldsymbol{\lambda} \equiv \mathbf{1}$ it is also both $\mathcal{D}$ - and $\mathcal{E}$ -nonvanishing. For any $D', D'' : V \rightarrow \mathbb{N}$ such that $\text{outdeg}(G) \leq D'$ and $\text{indeg}(G) \leq D''$, each variable z'_v may

be polarized $D'(v)$ times, and each variable z_v'' may be polarized $D''(v)$ times. The resulting polynomial

$$P^{D',D''}F(G; \{z'_{vi}\}, \{z''_{vj}\})$$

is still $\mathcal{H}$ -, $\mathcal{D}$ -, and $\mathcal{E}$ -nonvanishing, by repeated application of Proposition 10. From this point onward, the method of proof of Theorem 17 can be applied *mutatis mutandis*, and no new complications arise. Lacking a compelling application of the result, we leave the details to the reader.

REFERENCES

- [1] M. AISEN, I.J. SCHOENBERG, and A. WHITNEY, *On the generating functions of totally positive sequences*, Proc. Nat. Acad. Sci. U.S.A. **37** (1952), 303–307.
- [2] B.A. ASNER, JR., *On the total nonnegativity of the Hurwitz matrix*, SIAM J. Appl. Math. **18** (1970), 407–414.
- [3] Y.-B. CHOE, J.G. OXLEY, A.D. SOKAL and D.G. WAGNER, *Homogeneous multivariate polynomials with the half-plane property*, Adv. Appl. Math. **32** (2004), 88–187.
- [4] J. GARLOFF and D.G. WAGNER, *Hadamard products of stable polynomials are stable*, J. Math. Anal. Appl. **202** (1996), 797–809.
- [5] G.H. HARDY, J.E. LITTLEWOOD, and G.H. PÓLYA, “Inequalities”, Cambridge U.P. Cambridge, U.K., 1959.
- [6] O.J. HEILMANN and E.H. LIEB, *Theory of monomer-dimer systems*, Comm. Math. Phys. **25** (1972), 190–232.
- [7] S. KARLIN, “Total Positivity, vol. I”, Stanford U.P., Stanford, U.S.A., 1968.
- [8] J.H.B. KEMPERMAN, *A Hurwitz matrix is totally positive*, SIAM J. Math. Anal. **13** (1982), 331–341.
- [9] M. MARDEN, “Geometry of Polynomials”, Amer. Math. Soc., Providence, RI, 1966.
- [10] D. RUELLE, *Counting unbranched subgraphs*, J. Algebraic Combin. **9** (1999), 157–160.
- [11] D. RUELLE, *Zeros of graph-counting polynomials*, Comm. Math. Phys. **200** (1999), 43–56.
- [12] T. TAKAGI, *Note on the algebraic equations*, Proc. Phys.–Math. Soc. Japan **3** (1921), 175–179.
- [13] D.G. WAGNER, *Total positivity of Hadamard products*, J. Math. Anal. Appl. **163** (1992), 459–483.
- [14] D.G. WAGNER, *Multipartition series*, S.I.A.M. J. Discrete Math. **9** (1996), 529–544.
- [15] L. WEISNER, *Polynomials whose roots lie in a sector*, Amer. J. Math. **64** (1942), 55–60.

DEPARTMENT OF COMBINATORICS AND OPTIMIZATION, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO, CANADA N2L 3G1
 E-mail address: dgwagner@math.uwaterloo.ca